REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO N	OT RETURN YO	JR FORM TO T	HE ABOVE ORGANIZAT	ION.	,	
1. REPORT DA	ATE (DD-MM-YY	YY) 2. REP	ORT TYPE			3. DATES COVERED (From - To)
			Poster			12/04/2017
4. TITLE AND		**			5a. CON	NTRACT NUMBER
	Associated Acut	: Hyperkalemia	in a Swine Model of Sodi	ium Azide		
Toxicity					5b. GRA	ANT NUMBER
					SC. PRO	OGRAM ELEMENT NUMBER
6. AUTHOR(S)		##			5d. PRC	DJECT NUMBER
Maj Maddry, J	oseph					
					50 TAS	K NUMBER
					Se. IAS	R NOWIDER
					5f. WOF	RK UNIT NUMBER
7. PERFORMI	NG ORGANIZAT	ON NAME(S) A	ND ADDRESS(ES)			8. PERFORMING ORGANIZATION
	esearch Division		,			REPORT NUMBER
	Hall Loop, Bldg					
	i, TX 78236-990					17509
210-292-7141						
9. SPONSORII	NG/MONITORING	AGENCY NAN	/IE(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)
59th Clinical R	esearch Division					
1100 Willford	Hall Loop, Bldg	4430				
JBSA-Lackland	i, TX 78236-990	8				11. SPONSOR/MONITOR'S REPORT
210-292-7141						NUMBER(S)
	ION/AVAILABIL					
Approved for p	ublic release. Di	stribution is unli	mited.			
40 011001 511						
-	NTARY NOTES	whin Conforma	e, Falls Church, Virginia,	Dagamban 4. 2	017	
Science and Te	ciliology Leader	sinp Conterence	, rans Church, virginia,	December 4, 2	017	
14. ABSTRACT	•					
15. SUBJECT T	ERMS					
			T.2			
	CLASSIFICATIO		17. LIMITATION OF ABSTRACT	18. NUMBER OF		IE OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	C. IHIS PAGE		PAGES		Longoria
					19b. TEL	EPHONE NUMBER (Include area code)

Sodium Azide Associated Acute Hyperkalemia in a Swine Model of Sodium Azide Toxicit

PC Ng¹, JK Maddry¹,², VS Bebarta³, N Garrett⁴, S Boudreau², M Castaneda², K Canellis²

Wilford Hall Ambulatory Surgical Center, Lackland AFB, TX 3University of Colorado-Denver, 4Geneva Foundation, San Antonio, TX ¹Dept of Emergency Medicine, San Antonio Military Medical Center, San Antonio, TX, ²CREST Program,







Background

sodium azide poisoning. Virtually all of the used for sodium azide poisonings but have of those describe hypokalemia hours after extremely deadly. There is very little in the had limited success poisoning. Antidotes to cyanide have been information comes from case studies and each literature regarding the clinical course of Sodium azide (NaN3) poisonings are rare but

Objective

poisoning and develop novel treatments for To describe the clinical course of sodium azide

Methods

sustained for 1.5 minutes post apnea. Only sustained apnea 160 mg/mL produced consistent apnea but not doses at 10 mg/kg/min at concentrations of 1 minute by capnography. This rate was to 10 mg/kg/min until apnea was confirmed for mg/mL, was infused at doses ranging from 0.8 in concentrations ranging from 4 to 160 ventilate with an FiO2 of 0.21. Sodium azide, adjusted such that animals would spontaneous monitoring. After stabilization, anesthesia was femoral and pulmonary artery pressure intubated, and instrumented with continuous Twenty swine (45-55 kg) were anesthetized

Methods cont.

changes among groups over time. used to determine statistically significant Statistics: Repeated measures ANOVA was

Results

electrocardiogram and continued acidosis demonstrated elevated T waves on dextrose 50%, and calcium survived, but hyperkalemia died. Those treated with insulin, discontinued. Swine not treated for mmol/L) even after the infusion was mmol/L) and continued to rise (mean 7.7 significant hyperkalemia began at apnea (5.1 dose and concentration of NaN3 (n=14), hypotensive. In pigs infused with the highest pigs became hyperkalemic, acidotic and animals. Once the NaN3 infusion began, all mEq/L) and lactate (1.1 mmol/L) among the blood gases including potassium (mean 4.1 baseline vital signs, chemistries, or arterial (lactate mean 6.7 mmo/L). There were no significant differences in

	Baseline	Apnea	End of Study
PH	7.460	7.403	7.423
CO2	41.6	50.6	43.1
PO2	96.4	39.4	70.6
7	4.1	4.8	6.8
Ca2+	1.28	1.31	1.32
Lactate	1.1	1.5	10.5

Results Continued

Graph 1. Potassium trends over time in treated vs. non treated animals





Figure 1. ST elevation in animal #8611. NaN3 started at 0850 and off at 0853. Treated for hyperkalemia at 0852. ST changes noted at 0856.



_imitations

Animal model Infusion, not inhalation model

Conclusions

muscle breakdown. Model development is hyperkalemia is not caused by excessive exchange of potassium ions for hydrogen ions hyperkalemia is due, in part, to the intracellular hyperkalemia. We speculate that the findings in the animals demonstrate that in the face of metabolic acidosis. Pathology NaN3-poisoned swine acutely develop